

# Chronic Musculoskeletal Pain: Nonpharmacologic, Noninvasive Treatments

Diane M. Flynn, MD, MPH, Madigan Army Medical Center, Tacoma, Washington

Chronic low back pain, neck pain, hip and knee osteoarthritis, and fibromyalgia are the most common types of chronic musculoskeletal pain. Because no individual therapy has consistent benefit, a multimodal treatment approach to chronic musculoskeletal pain is recommended. Many nonpharmacologic, noninvasive treatment approaches yield small to moderate improvement and can be used with pharmacologic or more invasive modalities. Systematic reviews and guidelines support the effectiveness of various forms of exercise in improving pain and function in patients with chronic pain. Cognitive behavior therapy and mindfulness techniques appear to be effective for small to moderate short- and long-term improvement of chronic low back pain. Cognitive behavior therapy may also be effective for small short- and intermediate-term improvement of fibromyalgia. Spinal manipulation leads to a small benefit for chronic neck and low back pain. Acupuncture has a small to moderate benefit for low back pain and small benefit for nonpain fibromyalgia symptoms. Massage or myofascial release yields a small improvement in low back pain, hip and knee osteoarthritis, and fibromyalgia. Low reactive level laser therapy may provide short-term relief of chronic neck and low back pain, and ultrasound may provide short-term pain relief for knee osteoarthritis. Multidisciplinary rehabilitation may be effective for short- and at least intermediate-term improvement in pain and function for chronic low back pain and fibromyalgia. Patients should be encouraged to engage in a variety of therapies aligned with their preferences and motivation. (*Am Fam Physician*. 2020; 102(8):465-477. Copyright © 2020 American Academy of Family Physicians.)



Illustration by Jonathan Dimes

## A chronic musculoskeletal pain disorder

is the underlying diagnosis for 70% to 80% of those living with chronic pain.<sup>1</sup> Among the top 12 causes of disability in the United States, musculoskeletal disorders cause more than one-third of years lived with disability and are among the leading causes of disability worldwide.<sup>2,3</sup> Chronic low back pain, neck pain, hip and knee osteoarthritis, and fibromyalgia are the most common types of chronic musculoskeletal pain.

No one pain therapy alone consistently confers meaningful benefit for chronic musculoskeletal pain. On average, patients with chronic musculoskeletal pain experience small to moderate improvement in pain intensity and function from

any single therapy; therefore, a multimodal treatment approach that maximizes likelihood of benefit and minimizes risk of harm is recommended.<sup>4</sup>

A multimodal pain treatment plan (*Figure 1*) should emphasize self-management and may include the physical and psychological approaches summarized in this evidence review, as well as medications and herbal therapies, therapeutic injections, and surgery, which are beyond the scope of this article. Patients should be encouraged to engage in a variety of therapies aligned with their preferences and motivation.

## Patient Education

Patient education on neurophysiology, as part of a multicomponent self-management program, has demonstrated effectiveness for chronic low back pain.<sup>5</sup> Patient education may also reduce pain and nonpain symptoms in fibromyalgia.<sup>6</sup> Patient education on the diagnosis and on activity, lifestyle modification, and weight reduction,

**CME** This clinical content conforms to AAFP criteria for CME. See CME Quiz on page 461.

**Author disclosure:** No relevant financial affiliations.

if indicated, is recommended but not empirically proven for osteoarthritis.<sup>7</sup> No benefit has been established for chronic neck pain.<sup>8</sup> The evidence for patient education is summarized in *Table 1*.<sup>5-37</sup>

### Physical Activity

Physical activity that enhances or maintains muscle strength, physical fitness, flexibility, or overall health is recommended for most types of chronic pain.<sup>38</sup>

### CHRONIC LOW BACK PAIN

Research on exercise in general shows that in patients who have low back pain with or without lumbar radiculopathy, exercise leads to a small to moderate improvement in pain and a small improvement in function in the short term<sup>10-12</sup> (one to less than six months), small improvement in pain in the intermediate term (six to less than 12 months), and moderate improvement in pain in the long term (12 months or more).<sup>10</sup>

Motor control exercise is initially guided by a therapist, then continued independently, and focuses on improving coordination and function of the deep muscles that support the spine. It provides a moderate pain benefit and small functional benefit for low back pain that is slightly better than with general exercise for low back pain.<sup>11,12,23</sup>

Several movement disciplines improve pain and function. Yoga is recommended for chronic low back pain and results in small to moderate pain and functional benefits in the short and intermediate term.<sup>5,10-12,25-28</sup> Tai chi (a flowing series of movements that are coordinated with breathing) results in moderate pain benefit (equivalent to swimming) and small functional benefit for low back pain.<sup>10-12</sup>

### CHRONIC NECK PAIN

Although general exercise has no clear benefit for chronic neck pain, specific strengthening exercises of the neck, upper back, and shoulders reduce neck pain.<sup>10,14</sup> Combinations of muscle performance, muscle reeducation, mobility, and aerobic exercises improve neck pain and function.<sup>10</sup>

Motor control exercise provides moderate pain relief and small functional benefit for neck pain.<sup>24</sup> Evidence reviews of yoga for chronic neck pain have mixed results, with one review reporting a large decrease in pain and moderate improvement in function and another reporting insufficient evidence.<sup>25,29</sup> Tai chi leads to a small benefit for neck pain and function.<sup>25,33</sup>

## SORT: KEY RECOMMENDATIONS FOR PRACTICE

Clinical recommendation	Evidence rating*
Regular exercise is recommended for patients with chronic musculoskeletal pain. <sup>10-12,18-23</sup> Because no specific type of exercise is clearly superior, patients should be encouraged to engage in the type of low-impact exercise they prefer.	<b>B</b>
Encourage yoga for patients with chronic low back pain, <sup>10,11,25,27,28</sup> lumbar radiculopathy, <sup>11</sup> knee osteoarthritis, <sup>30,31</sup> or fibromyalgia. <sup>30,32</sup>	<b>B</b>
Encourage cognitive behavior therapy for patients with chronic low back pain <sup>10-12,25</sup> or fibromyalgia. <sup>10,41,42</sup>	<b>B</b>
Encourage mindfulness-based stress reduction for patients with chronic low back pain <sup>10-12,25</sup> or fibromyalgia. <sup>10</sup>	<b>B</b>
Consider spinal manipulation for patients with chronic low back pain <sup>10</sup> or neck pain. <sup>48,49</sup>	<b>B</b>
Consider acupuncture for patients with chronic low back pain, <sup>10-12,52</sup> neck pain, <sup>10,52</sup> or fibromyalgia. <sup>10,54</sup>	<b>B</b>
Consider massage or myofascial release for low back pain, <sup>10,58,59</sup> neck pain, <sup>10</sup> hip <sup>10</sup> and knee <sup>60,61</sup> osteoarthritis, and fibromyalgia. <sup>10</sup>	<b>B</b>
Consider multi- or interdisciplinary rehabilitation for patients with chronic low back pain or fibromyalgia that does not respond to initial therapies, or who have a significant psychological component. <sup>10-12,71</sup>	<b>B</b>

**A** = consistent, good-quality patient-oriented evidence; **B** = inconsistent or limited-quality patient-oriented evidence; **C** = consensus, disease-oriented evidence, usual practice, expert opinion, or case series. For information about the SORT evidence rating system, go to <https://www.aafp.org/afpsort>.


\*—All are based on multiple consistent systematic reviews, including Cochrane reviews, of low- to moderate-quality studies and evidence-based guidelines.

### HIP AND KNEE OSTEOARTHRITIS

Exercise improves symptoms of hip and knee osteoarthritis.<sup>7,10</sup> Clinical trials of knee osteoarthritis have studied walking, gait training, mobility exercise, and exercise combinations. Trials of hip osteoarthritis have included strength training, stretching, neuromuscular control exercises, and endurance training. For knee osteoarthritis, exercise results in small to moderate improvements in pain and function in the short and intermediate term, as well as small long-term improvements in function but not pain.<sup>10,16</sup> For hip osteoarthritis, exercise leads to small short-term improvements in pain and function and small intermediate-term improvements in function but not pain.<sup>10</sup>

Aquatic exercise results in small short-term improvements in pain, function, and health-related quality of life for hip and knee osteoarthritis.<sup>18</sup> Yoga results in improved

FIGURE 1



**Note:** Treatments that overlap with self-management have important components of patient engagement.

Potential components of a multimodal pain treatment plan.

pain, mobility, and function for knee osteoarthritis.<sup>30,31</sup> There is mixed evidence for tai chi in the treatment of knee osteoarthritis.<sup>10,34</sup>

### FIBROMYALGIA

Several systematic reviews support exercise for patients with fibromyalgia. General exercise results in small improvements in pain and function in the short term, small improvements in function but not pain in the intermediate term, and no improvement in pain or function in the long term.<sup>10,19</sup>

Clinical trials of specific forms of exercise for fibromyalgia show that resistance exercise such as lifting weights, aerobic exercise such as walking, aquatic exercise, and various combinations of exercise all yield small improvements in pain and function, and some exercise types are beneficial for other fibromyalgia symptoms.<sup>19-22</sup> Yoga improves pain, fatigue, depression, health-related quality of life, and function in patients with fibromyalgia.<sup>30,32</sup> Tai chi or qigong (many single movements, each of which may be repeated to address a specific concern) improves short-term pain and function.<sup>10</sup> Tai chi also improves other common fibromyalgia symptoms.<sup>32</sup>

### SUMMARY

Regular engagement in physical activity through exercise, physical therapy, and/or movement disciplines is among the most consistently beneficial approaches to the treatment of chronic musculoskeletal pain and should be the foundation of a multimodal treatment plan. Because no specific type of exercise is clearly superior, patients should be encouraged to engage in the type of low-impact exercise they prefer. The evidence for physical activity is summarized in *Table 1*.<sup>5-37</sup>

### Psychological Therapies, Mindfulness, and Muscle Relaxation

#### LOW BACK PAIN

Research on psychological therapies for chronic musculoskeletal pain has focused primarily on cognitive behavior therapy (CBT) and confirms its benefit for low back pain, including small to moderate short-, intermediate-, and long-term improvement in pain. In addition, CBT confers small functional improvement in the short, intermediate, and long term.<sup>10-12,25</sup> The most recent reviews of CBT also report improvement in function for up to two years.<sup>10,25</sup> Operant therapy, which focuses on eliminating

TABLE 1

## Patient Education and Physical Activity for Chronic Musculoskeletal Pain: Review of the Evidence

Intervention	Common types of pain (SOR rating*)		
	Low back	Neck	Radicular/referred
Patient education	<ul style="list-style-type: none"> <li>● Education on neurophysiology is recommended as part of a multicomponent self-management program (C)<sup>5</sup></li> <li>● Insufficient evidence for "back schools"<sup>9</sup></li> </ul>	<ul style="list-style-type: none"> <li>● Insufficient evidence<sup>8</sup></li> </ul>	No studies found based on search criteria
Exercise	<ul style="list-style-type: none"> <li>● General exercise Overall benefit (C)<sup>5</sup> Short term: ↓ to ↓↓ pain and ↑ function (A)<sup>10-12</sup> Intermediate term: ↓ pain, no effect on function (B)<sup>10</sup> Long term: ↓↓ pain, no effect on function (B)<sup>10</sup></li> <li>● Giving patients a pedometer, walking diary, and instructions to walk at least four days per week ≈ physical therapy or group exercise for ↓ pain and ↑ function (B)<sup>13</sup></li> <li>● Pilates is recommended but provides no additional pain or functional benefit when added to general exercise (C)<sup>5,11</sup></li> </ul>	<ul style="list-style-type: none"> <li>● General exercise: no short-term benefit and insufficient evidence of long-term benefit<sup>10</sup></li> <li>● Strengthening exercises of the neck, upper back, and shoulders: ↓ to ↓↓ pain (B)<sup>14</sup></li> </ul>	<ul style="list-style-type: none"> <li>● General exercise: ↓ pain and ↑ function for lumbar radiculopathy (C)<sup>11</sup></li> <li>● Pilates: insufficient evidence for lumbar radiculopathy<sup>11</sup></li> </ul>
Physical or occupational therapy	<ul style="list-style-type: none"> <li>● Clinician-directed exercise is recommended (C)<sup>5</sup></li> <li>● Motor control exercises: ↓↓ pain and ↑ function (B)<sup>11,12,23</sup></li> <li>● Motor control exercise is slightly better than general exercise (B)<sup>11,12,23</sup></li> </ul>	<ul style="list-style-type: none"> <li>● Mixed exercise† Short term: ↓ pain and ↑ function (B)<sup>10</sup> Long term: ↑ function (B)<sup>10</sup></li> <li>● Motor control exercises: ↓↓ pain and ↑ function (B)<sup>24</sup></li> </ul>	No studies found based on search criteria
Yoga	<ul style="list-style-type: none"> <li>● General benefit (C)<sup>5</sup></li> <li>● Short term: ↓ to ↓↓ pain and ↑ to ↑↑ function (B)<sup>10-12,25-28</sup></li> <li>● Intermediate term: ↓ to ↓↓ pain and ↑ to ↑↑ function (B)<sup>10,27</sup></li> <li>● Long term: interpretation differs (limited benefit vs. no benefit)<sup>10,27,28</sup></li> <li>● Slightly better for pain than other exercise (B)<sup>10,11,27</sup></li> </ul>	<ul style="list-style-type: none"> <li>● Interpretation differs: ↓↓ pain and ↑↑ function vs. insufficient evidence<sup>25,29</sup></li> </ul>	<ul style="list-style-type: none"> <li>● ↓ pain and ↑ function for lumbar radiculopathy (C)<sup>11</sup></li> </ul>
Tai chi	<ul style="list-style-type: none"> <li>● ↓↓ pain and ↑ function (B)<sup>10-12</sup></li> <li>● Tai chi ≈ swimming but superior to backward walking or jogging for pain (B)<sup>11</sup></li> </ul>	<ul style="list-style-type: none"> <li>● ↓ pain and ↑ function (B)<sup>25,33</sup></li> <li>● Tai chi ≈ conventional neck exercises (B)<sup>25,33</sup></li> </ul>	No studies found based on search criteria
Qigong¶	<ul style="list-style-type: none"> <li>● Exercise is slightly better for pain in the short term and for function in the intermediate term<sup>10</sup></li> </ul>	<ul style="list-style-type: none"> <li>● Interpretation differs: small to no benefit<sup>35-37</sup></li> </ul>	No studies found based on search criteria
Alexander technique**	No studies found based on search criteria	<ul style="list-style-type: none"> <li>● Short and intermediate term: ↑ function (B); insufficient evidence for pain<sup>10</sup></li> <li>● Long term: insufficient evidence<sup>10</sup></li> </ul>	No studies found based on search criteria

**Note:** short term = 1 to < 6 months; intermediate term = 6 to < 12 months; long term = ≥ 12 months; ↓ = small or undefined decrease; ↑ small or undefined increase; ↓↓ moderate decrease; ↑↑ moderate increase; ↓↓↓ = large decrease; small = 5 to 10 points on a 100-point scale or SMD of 0.2 to < 0.5; moderate = 10 to 20 points on a 100-point scale or SMD of 0.5 to < 0.8; large = ≥ 20 points on a 100-point scale or SMD ≥ 0.8; ● = evidence to support; ● = insufficient or inconsistent evidence; ● = evidence against; ● = comparative effectiveness with other therapies.

HRQL = health-related quality of life; SMD = standardized mean difference; SOR = strength of recommendation.

\*—A = consistent, good-quality patient-oriented evidence; B = inconsistent or limited-quality patient-oriented evidence; C = consensus, disease-oriented evidence, usual practice, expert opinion, or case series. For information about the SORT evidence rating system, go to <https://www.aafp.org/afpsort>.

†—Mixed exercise: includes a combination of exercises. For neck pain, it must include at least three of the following types: muscle performance, mobility, muscle reeducation, and aerobic. For fibromyalgia, it must include at least two of the following types: flexibility, motor, or aerobic.

## Osteoarthritis

- Education on the diagnosis and on activity, lifestyle modification, and weight reduction, if indicated, is recommended; effectiveness has not been empirically proven (C)<sup>7</sup>

### Hip

#### General exercise

- Short term: ↓ pain and ↑ function (B)<sup>10</sup>
- Intermediate term: ↑ function, no pain benefit (B)<sup>10</sup>
- Long term: insufficient evidence<sup>10</sup>

### Knee

#### General exercise

Overall benefit (C)<sup>15</sup>

Short term: ↓ to ↓↓ pain and ↑ to ↑↑ function (B)<sup>10,16</sup>

Intermediate term: ↓↓ pain and ↑ to ↑↑ function (B)<sup>10,16</sup>

Long term: ↑ function, no pain benefit (B)<sup>10</sup>

- Aerobic exercise: ↓ pain and ↑ function (C)<sup>17</sup>
- Strengthening exercise: ↓ pain and ↑ function (C)<sup>15</sup>
- Aquatic exercise: short-term ↓ pain, ↑ function, and ↑ HRQL (B)<sup>18</sup>

- Therapeutic exercise focusing on regaining quadriceps and gluteal strength, flexibility, range of motion, and aerobic condition: ↓ pain and ↑ function (C)<sup>7</sup>
- Manual physical therapy when added to traditional physical therapy and supervised exercise may further ↓ pain and ↑ function (C)<sup>7</sup>

- Knee: ↓ pain (C), ↑ mobility (B), and ↑ function (B)<sup>30,31</sup>

- Interpretation differs: limited benefit vs. insufficient evidence<sup>10,34</sup>

No studies found based on search criteria

No studies found based on search criteria

## Fibromyalgia

- ↓ pain, ↓ anxiety, ↓ perception of illness, ↓ catastrophizing (C)<sup>6</sup>

### General exercise

- Short term: ↓ pain and ↑ function (B)<sup>10,19</sup>
- Intermediate term: ↑ function, no pain benefit (B)<sup>10</sup>
- Long term: no pain or functional benefit (B)<sup>10</sup>
- Mixed exercise†  
Short term: ↓ pain, ↑ function, ↑ HRQL, ↓ fatigue, and ↓ stiffness  
Long term: ↑ HRQL, ↓ fatigue, and ↑ function (B)<sup>20</sup>
- Aerobic exercise: ↑ HRQL, ↓ pain, and ↑ function, but no benefit for stiffness or fatigue (B)<sup>21</sup>
- Resistance exercise‡: ↓ pain and ↑ function (B)<sup>19</sup>
- Aquatic exercise: ↓↓ stiffness, ↓ pain, ↑ function, and ↑ wellness (B)<sup>22</sup>

No studies found based on search criteria

- ↓↓ pain and ↑ function; improves fatigue, depression, and HRQL (B)<sup>30,32</sup>

- Short term: ↓↓ pain and ↑ function; improves sleep, fatigue, mood, and HRQL (B)<sup>10,32</sup>

- Intermediate and long term: insufficient evidence<sup>10</sup>

- Short term: ↓↓ pain and ↑ function (B)<sup>10</sup>

- Intermediate and long term: insufficient evidence<sup>10</sup>

No studies found based on search criteria

‡—Resistance exercise is the use of weights, bands, etc., to provide resistance to movement, which builds muscle strength and endurance.

§—Motor control exercise is initially guided by a therapist, then continued independently, and focuses on improving coordination and function of the deep muscles that support the spine.

||—Tai chi is a flowing series of movements that are coordinated with breathing.

¶—Qigong includes many single movements, each of which may be repeated to address a specific concern.

\*\*—The Alexander technique uses habitual patterns of posture and movement to reduce muscle tension. A video of the technique is available at <https://www.youtube.com/watch?v=niFdH63McSA>.

harmful pain behaviors and increasing beneficial behaviors by using the brain’s reward system, has a small benefit for low back pain but not function.<sup>11</sup> Evidence is insufficient regarding the effects of psychological therapies for lumbar radiculopathy.<sup>11</sup>

Mindfulness-based stress reduction (the daily practice of focusing attention on the present moment in a non-judgmental way) results in small to moderate short-term improvement and small intermediate-term improvement in

low back pain.<sup>10-12,25</sup> Findings differ regarding its impact on function and long-term outcomes.<sup>10-12,25</sup> One trial comparing mindfulness-based stress reduction or CBT with usual care for low back pain found that both approaches had moderate benefit for pain and function, with improvements roughly equivalent after one year; the benefit persisted after two years but was not statistically significant.<sup>39</sup>

Progressive muscle relaxation (reducing muscle tension by intentionally tensing, then relaxing, successive muscle

TABLE 2

**Psychological Therapies and Mindfulness Techniques for Chronic Musculoskeletal Pain: Review of the Evidence**

Intervention	Common types of pain (SOR rating*)		
	Low back	Neck	Radicular/referred
Psychological therapies (primarily CBT)	<ul style="list-style-type: none"> <li>● CBT</li> <li>Short and intermediate term: ↓ to ↓↓ pain and ↑ function (B)<sup>10-12,25</sup></li> <li>Long term: ↓ pain and ↑ function (B)<sup>10,25</sup></li> <li>● Operant therapy†: ↓ pain, no effect on function (B)<sup>11</sup></li> <li>● CBT ≈ exercise (B)<sup>11,12</sup></li> <li>● No clear benefit of one psychological therapy over another (B)<sup>11,12</sup></li> </ul>	<ul style="list-style-type: none"> <li>● Short term: insufficient evidence<sup>25,40</sup></li> <li>● CBT ≈ other treatments (medications, patient education, physical therapy, manual therapy, exercise) in ↓ pain and ↓ disability (B)<sup>40</sup></li> </ul>	<ul style="list-style-type: none"> <li>● Insufficient evidence for lumbar radiculopathy<sup>11</sup></li> </ul>
Electromyography biofeedback	<ul style="list-style-type: none"> <li>● ↓↓ pain, no effect on function (B)<sup>11,12</sup></li> </ul>	No studies found based on search criteria	No studies found based on search criteria
Mindfulness-based stress reduction‡	<ul style="list-style-type: none"> <li>● Short-term pain: ↓ to ↓↓ (B)<sup>5,10-12,25</sup></li> <li>● Intermediate-term pain: ↓ (B)</li> <li>● Short-term function: inconsistent evidence<sup>5,10</sup></li> <li>● Long-term pain and function: inconsistent evidence<sup>10,25</sup></li> <li>● CBT ≈ mindfulness-based stress reduction in decreasing pain and improving function at one year, which was partially sustained but no longer statistically significant at two years (B)<sup>39</sup></li> </ul>	<ul style="list-style-type: none"> <li>● Insufficient evidence<sup>25</sup></li> </ul>	No studies found based on search criteria
Other meditative therapies	<ul style="list-style-type: none"> <li>● Progressive muscle relaxation§: ↓↓ pain and ↑↑ function (B)<sup>11,12</sup></li> </ul>	<ul style="list-style-type: none"> <li>● Physical therapy–led relaxation training has no benefit (B)<sup>10</sup></li> <li>● Progressive muscle relaxation ≈ cupping   for ↓ pain (B)<sup>44</sup></li> </ul>	No studies found based on search criteria

**Note:** short term = 1 to < 6 months; intermediate term = 6 to < 12 months; long term = ≥ 12 months; ↓ = small or undefined decrease; ↑ small or undefined increase; ↓↓ moderate decrease; ↑↑ moderate increase; small = 5 to 10 points on a 100-point scale or SMD of 0.2 to < 0.5; moderate = 10 to 20 points on a 100-point scale or SMD of 0.5 to < 0.8; ● = evidence to support; ● = insufficient or inconsistent evidence; ● = evidence against; ● = comparative effectiveness with other therapies.

CBT = cognitive behavior therapy; SMD = standardized mean difference; SOR = strength of recommendation.

\*—A = consistent, good-quality patient-oriented evidence; B = inconsistent or limited-quality patient-oriented evidence; C = consensus, disease-oriented evidence, usual practice, expert opinion, or case series. For information about the SORT evidence rating system, go to <https://www.aafp.org/afpsort>.



groups) results in moderate improvement of low back pain and function.<sup>11,12</sup> Low-quality evidence suggests that electromyography biofeedback leads to a moderate reduction in low back pain.<sup>11,12</sup>

**NECK PAIN**

Findings on the effects of CBT for neck pain are limited because of the absence of follow-up beyond the active treatment period.<sup>25,40</sup>

**OSTEOARTHRITIS AND FIBROMYALGIA**

Psychological therapies yield small short-term improvement of knee osteoarthritis pain but not function.<sup>10</sup> For fibromyalgia, CBT has a small short-term benefit for pain, mood, and disability and intermediate-term benefit for function. Interpretation of evidence differs regarding short-term functional impact and intermediate-term pain impact.<sup>10,41,42</sup> One trial found that, when compared with pharmacologic therapies, CBT provided equivalent pain relief and superior functional benefit after six months in patients with fibromyalgia.<sup>10</sup> Mindfulness-based stress reduction leads to small improvements in pain and function for fibromyalgia in the intermediate term.<sup>10,43</sup> Biofeedback has no clear benefit for fibromyalgia.<sup>41</sup>

**SUMMARY**

CBT and mindfulness-based stress reduction appear to be effective for small to moderate short- and long-term improvement of low back pain. Biofeedback and progressive muscle relaxation may also be short-term treatment options for chronic low back pain. CBT appears to be effective for small short-term improvement of knee osteoarthritis pain and short- and intermediate-term improvement of fibromyalgia. Mindfulness-based stress reduction improves intermediate-term pain and function in fibromyalgia. The evidence for these techniques is summarized in *Table 2*.<sup>5,10-12,25,39-42,44-46</sup>

**Manual Therapies and Acupuncture**

Spinal manipulation (osteopathic or chiropractic) for low back pain leads to small short-term improvements in function, and inconsistent reports show small to no improvement in short-term pain intensity.<sup>10-12,47</sup> Spinal manipulation results in small improvements in low back pain and function in the intermediate term, but there is insufficient evidence regarding long-term effects.<sup>10</sup> Evidence on the effectiveness of spinal manipulation for lumbar radiculopathy is mixed.<sup>11,12</sup> Spinal manipulation improves chronic neck pain and function, but interpretation of evidence differs on whether thoracic is superior to cervical manipulation.<sup>48-50</sup> Evidence is insufficient regarding the effectiveness of spinal manipulation for fibromyalgia.<sup>51</sup>

Acupuncture provides small to moderate short-term and small long-term, but not intermediate-term, benefit for low back pain when compared with sham acupuncture.<sup>10-12,52</sup> Overall,

Knee osteoarthritis	Fibromyalgia
<ul style="list-style-type: none"> <li>Short term: ↓ pain, no functional benefit (B)<sup>10</sup></li> </ul>	<ul style="list-style-type: none"> <li>Short term: ↓ pain, ↑ mood, and ↓ disability (B)</li> <li>Short-term function: interpretation differs<sup>10,41,42</sup></li> <li>Intermediate term: ↑ to ↑↑ function (B)<sup>10,41,42</sup></li> <li>Intermediate-term pain: interpretation of the impact on pain differs (↓ pain vs. no benefit)<sup>10,41,42</sup></li> <li>Long term: insufficient evidence<sup>10</sup></li> <li>CBT ≈ pregabalin (Lyrica) or duloxetine (Cymbalta) for pain relief; CBT better for function after six months (B)<sup>10,42</sup></li> </ul>
No studies found based on search criteria	<ul style="list-style-type: none"> <li>Little or no improvement in function, pain, mood (B)<sup>41</sup></li> </ul>
No studies found based on search criteria	<ul style="list-style-type: none"> <li>Short term: no benefit (B)<sup>10</sup></li> <li>Intermediate term: ↓ pain and ↑ function (B)<sup>10</sup></li> <li>Long term: insufficient evidence<sup>10</sup></li> </ul>
No studies found based on search criteria	<ul style="list-style-type: none"> <li>Guided imagery/hypnosis: ↓ pain (B)<sup>45</sup></li> <li>Progressive muscle relaxation: insufficient evidence<sup>46</sup></li> </ul>

†—Operant therapy focuses on eliminating harmful pain behaviors and increasing beneficial behaviors by leveraging the brain’s reward system.  
 ‡—Mindfulness-based stress reduction is the daily practice of focusing attention on the present moment in a nonjudgmental way.  
 §—Progressive muscle relaxation is a method of reducing muscle tension by intentionally tensing, then relaxing, successive muscle groups.  
 ||—Cupping is the application of suction to the skin through the use of glass, ceramic, bamboo, or plastic cups.

*Information from references 5, 10-12, 25, 39-42, and 44-46.*

the quality of studies on acupuncture is low. Impact on function varies from small short-term benefit to no benefit.<sup>10-12</sup> Evidence is insufficient regarding the effectiveness of acupuncture for lumbar radicular pain.<sup>11</sup> Acupuncture improves function for chronic neck pain, but the effect on pain is inconsistent.<sup>10,52</sup> The effect of acupuncture on knee osteoarthritis varies, with some studies finding a small benefit for pain in the short and long term and others finding no benefit.<sup>10,52</sup> Acupuncture has no clear benefit for hip osteoarthritis.<sup>53</sup> For fibromyalgia, acupuncture improves function and

stiffness but has no effect on pain.<sup>10,54</sup> Dry needling (insertion of acupuncture needles into myofascial trigger points) provides short-term relief for low back and neck pain.<sup>55-57</sup>

Massage leads to short-term improvement in low back pain and function; there is no pain benefit, conflicting interpretation of functional impact at intermediate-term follow-up, and insufficient evidence of long-term benefit.<sup>10,58,59</sup> Compared with exercise, massage provides no additional benefit for pain or function in the intermediate term.<sup>10</sup> Evidence is insufficient regarding the effectiveness

TABLE 3

Manual Therapies and Acupuncture for Chronic Musculoskeletal Pain: Review of the Evidence

Common types of pain (SOR rating\*)

Intervention	Low back	Neck	Radicular/referred
Spinal manipulation (osteopathic or chiropractic)	<ul style="list-style-type: none"> <li>● Short-term function: ↑ (B)<sup>10</sup></li> <li>● Short-term pain: inconsistent reports (small to no benefit)<sup>10-12,47</sup></li> <li>● Intermediate term: ↓ pain and ↑ function (B)<sup>10</sup></li> <li>● Long term: insufficient evidence<sup>10</sup></li> <li>● Spinal manipulation ≈ exercise for short- and intermediate-term ↓ pain and ↑ function (B)<sup>10</sup></li> </ul>	<ul style="list-style-type: none"> <li>● Various types of spinal manipulation and/or mobilization: ↓ pain and ↑ function (B)<sup>48</sup></li> <li>● Interpretation differs on the relative benefit of thoracic vs. cervical manipulation (B)<sup>49,50</sup></li> </ul>	<ul style="list-style-type: none"> <li>● Lumbar radicular pain: interpretation differs (↓ back and leg pain when added to home exercise and advice vs. insufficient evidence)<sup>11,12</sup></li> </ul>
Acupuncture (compared with sham acupuncture)	<ul style="list-style-type: none"> <li>● Short-term pain: ↓ to ↓↓ (B)<sup>10-12,52</sup></li> <li>● Short-term function: interpretation differs (small to no benefit)<sup>11,12</sup></li> <li>● Intermediate term: no benefit (B)<sup>10</sup></li> <li>● Long term: ↓ pain (B)<sup>10,52</sup></li> </ul>	<ul style="list-style-type: none"> <li>● Short and intermediate term: ↑ function<sup>10</sup></li> <li>● Pain: interpretation differs (small benefit vs. insufficient evidence)<sup>10,52</sup></li> <li>● Long term: no benefit (B)<sup>10</sup></li> </ul>	<ul style="list-style-type: none"> <li>● Lumbar radicular pain: insufficient evidence<sup>11</sup></li> </ul>
Dry needling†	<ul style="list-style-type: none"> <li>● Short term: ↓ pain (B)<sup>55</sup></li> <li>● Dry needling is better than sham therapy, physical therapy, and local anesthetic injection for ↓ pain and ↑ function (B)<sup>63</sup></li> </ul>	<ul style="list-style-type: none"> <li>● Short term (upper trapezius trigger points): ↓ pain (B)<sup>56,57</sup></li> </ul>	<ul style="list-style-type: none"> <li>● No studies found based on search criteria</li> </ul>
Manual therapies‡	<p><b>Massage</b></p> <ul style="list-style-type: none"> <li>● Short term: ↓ to ↓↓ pain and ↑ function (B)<sup>10,58,59</sup></li> <li>● Intermediate term: interpretation differs (↑ function vs. no benefit)<sup>10,59</sup></li> <li>● Long term: insufficient evidence<sup>10</sup></li> <li>● Massage ≈ exercise for intermediate-term pain (B)<sup>10</sup></li> </ul>	<ul style="list-style-type: none"> <li>● Short term: ↓↓ pain and ↑ function (B)<sup>10</sup></li> <li>● Intermediate term: no benefit on function and no evidence for pain (B)<sup>10</sup></li> <li>● Long term: insufficient evidence<sup>10</sup></li> </ul>	<ul style="list-style-type: none"> <li>● Insufficient evidence for lumbar radiculopathy<sup>11</sup></li> </ul>

**Note:** short term = 1 to < 6 months; intermediate term = 6 to < 12 months; long term = ≥ 12 months; ↓ = small or undefined decrease; ↑ = small or undefined increase; ↓↓ moderate decrease; ↑↑ moderate increase; small = 5 to 10 points on a 100-point scale or SMD of 0.2 to < 0.5; moderate = 10 to 20 points on a 100-point scale or SMD of 0.5 to < 0.8; ● = evidence to support; ● = insufficient or inconsistent evidence; ● = evidence against; ● = comparative effectiveness with other therapies.

SMD = standardized mean difference; SOR = strength of recommendation.

Information from references 10-12, 26, and 47-63.



of massage for lumbar radiculopathy.<sup>11</sup> For neck pain, massage provides moderate pain relief and small functional improvement in the short term.<sup>10</sup>

Massage for knee osteoarthritis provides short-term benefit for pain and function.<sup>60</sup> Other manual therapies for knee osteoarthritis provide moderate benefit for function but have no impact on pain.<sup>61</sup> For hip osteoarthritis, manual therapies provide a small short-term improvement in pain and function and a small intermediate-term improvement in function.<sup>10</sup> Evidence regarding the short-term impact of

massage for fibromyalgia is mixed.<sup>26,62</sup> Myofascial release for fibromyalgia yields a small intermediate-term increase in function and a small long-term reduction in pain.<sup>10</sup>

**SUMMARY**

Among manual therapies and acupuncture, mostly limited to short-term results, spinal manipulation is modestly effective for chronic low back and neck pain; acupuncture may be modestly to moderately effective for low back pain and nonpain symptoms of fibromyalgia and modestly or inconsistently effective for neck pain; and massage and/or myofascial release may be moderately effective for neck pain and modestly effective for low back pain, hip and knee osteoarthritis, and fibromyalgia. The evidence for manual therapies and acupuncture is summarized in *Table 3*.<sup>10-12,26,47-63</sup>

**Treatment Devices**

There is insufficient evidence for the effectiveness of transcutaneous electrical nerve stimulation in the treatment of chronic low back pain with or without associated lumbar radiculopathy, neck pain, knee osteoarthritis, or fibromyalgia.<sup>10,11,64-66</sup> There is insufficient evidence for the use of lumbar supports for low back pain or braces for knee osteoarthritis.<sup>5,11,67</sup> There is no short-term benefit of traction for low back pain with or without radiculopathy and insufficient evidence of benefit for chronic neck pain.<sup>10,11,68</sup>

For low back pain, low reactive level laser therapy (LLLT; the application of light in wavelengths that penetrate the skin, triggering biochemical changes to reduce pain and inflammation) provides a small to moderate short-term improvement in pain and a small improvement in function, based on low-quality evidence.<sup>10,11</sup> For chronic neck pain, moderate evidence shows that LLLT provides moderate short-term benefit for pain and function.<sup>10</sup> LLLT may provide short-term relief of pain and nonpain fibromyalgia symptoms.<sup>69</sup> There is insufficient evidence for the benefit of LLLT in the treatment of lumbar radiculopathy or knee osteoarthritis.<sup>10,11</sup> Ultrasound is not effective for low back pain or knee osteoarthritis.<sup>10,11</sup>

**SUMMARY**

LLLT may have short-term benefits for low back pain, neck pain, and fibromyalgia. Available evidence does not support use of other devices, including transcutaneous electrical

Osteoarthritis	Fibromyalgia
No studies found based on search criteria	● Insufficient evidence <sup>51</sup>
<p><b>Hip</b></p> <p>● No clear benefit (B)<sup>55</sup></p> <p><b>Knee</b></p> <p>● Interpretation differs: ↓ pain in short and long term vs. no pain benefit<sup>10,52</sup></p>	<p>● Short and intermediate term: ↑ function and ↓ stiffness, but no pain benefit (B)<sup>10,54</sup></p> <p>● Long term: insufficient evidence<sup>10</sup></p>
No studies found based on search criteria	No studies found based on search criteria
<p><b>Hip</b></p> <p>● Manual therapies: short-term ↓ pain and ↑ function and intermediate-term ↑ function (B)<sup>10</sup></p> <p><b>Knee</b></p> <p>● Massage: short-term ↓ pain and ↑ function (B)<sup>60</sup></p> <p>● Other manual therapies: ↑↑ function, no pain benefit (B)<sup>61</sup></p>	<p>● Massage: interpretation differs on short-term impact (↓ pain, ↓ anxiety, and ↓ depression to no benefit)<sup>26,62</sup></p> <p>● Myofascial release</p> <p>Intermediate term: ↑ function (B)<sup>10</sup></p> <p>Long term: ↓ pain but no impact on function (B)<sup>10</sup></p>

\*—A = consistent, good-quality patient-oriented evidence; B = inconsistent or limited-quality patient-oriented evidence; C = consensus, disease-oriented evidence, usual practice, expert opinion, or case series. For information about the SORT evidence rating system, go to <https://www.aafp.org/afpsort>.

†—Dry needling is the insertion of acupuncture needles into myofascial trigger points.

‡—Manual therapies are used by physical therapists and include manipulation, mobilization, and massage.

nerve stimulation, traction, lumbar supports, and ultrasound. The evidence for treatment devices is summarized in *Table 4*.<sup>5,10-12,64-69</sup>

**Multi- or Interdisciplinary Pain Rehabilitation**

Multi- or interdisciplinary pain rehabilitation includes coordinated care by a team comprised of combinations of

clinicians, physical or occupational therapists, and behavior therapists.<sup>70</sup> It is recommended for patients with low back pain that does not respond sufficiently to first-line therapies. Multi- or interdisciplinary pain rehabilitation provides small to moderate short-term improvement in pain, function, and disability and small intermediate-term benefit for pain and function.<sup>10-12,71</sup> Assessments of long-term benefit

TABLE 4

**Treatment Devices for Chronic Musculoskeletal Pain: Review of the Evidence**

Intervention	Common types of pain (SOR rating*)				
	Low back	Neck	Radicular/referred	Knee osteoarthritis	Fibromyalgia
Transcutaneous electrical nerve stimulation	● Interpretation differs (ineffective vs. insufficient evidence) <sup>64</sup>	● Insufficient evidence for chronic neck pain <sup>65</sup>	● Insufficient evidence for lumbar radiculopathy <sup>11</sup>	● Short or long term: insufficient evidence <sup>10</sup> ● Intermediate term: no pain or functional benefit (B) <sup>10</sup>	● Insufficient evidence <sup>66</sup>
Braces/supports	● Lumbar supports: insufficient evidence <sup>5,11</sup>	No studies found based on search criteria	No studies found based on search criteria	● Knee braces and foot/ankle orthoses: insufficient evidence <sup>67</sup>	No studies found based on search criteria
Traction	● No short-term benefit (B) <sup>10</sup>	● Insufficient evidence <sup>68</sup>	● No effect for lumbar radiculopathy (C) <sup>11</sup>	No studies found based on search criteria	No studies found based on search criteria
Low reactive level laser therapy†	● Short term: ↓ to ↓↓ pain and ↑ function (B) <sup>10-12</sup> ● Intermediate term: no benefit (B) <sup>10</sup> ● Long term: insufficient evidence <sup>10</sup>	● Short term: ↓↓ pain and ↑↑ function (B) <sup>10</sup> ● Intermediate and long term: insufficient evidence <sup>10</sup>	● Insufficient evidence for lumbar radiculopathy <sup>11</sup>	● Insufficient evidence <sup>10</sup>	● Short term: ↓ pain, ↓ Fibromyalgia Impact Questionnaire score, ↓ fatigue, ↓ stiffness, ↑ mood <sup>69</sup>
Ultrasound	● Short-term pain: no benefit (B) <sup>10</sup> ● Short-term function: insufficient evidence ● Intermediate and long term: insufficient evidence <sup>10</sup>	No studies found based on search criteria	No studies found based on search criteria	● Short- and intermediate-term pain and function: no benefit (B) <sup>10</sup> ● Long term: insufficient evidence <sup>10</sup>	No studies found based on search criteria

**Note:** short term = 1 to < 6 months; intermediate term = 6 to < 12 months; long term = ≥ 12 months; ↓ = small or undefined decrease; ↑ small or undefined increase; ↓↓ moderate decrease; ↑↑ moderate increase; small = 5 to 10 points on a 100-point scale or SMD of 0.2 to < 0.5; moderate = 10 to 20 points on a 100-point scale or SMD of 0.5 to < 0.8; ● = evidence to support; ● = insufficient or inconsistent evidence; ● = evidence against; ● = comparative effectiveness with other therapies.

SMD = standardized mean difference; SOR = strength of recommendation.

\*—A = consistent, good-quality patient-oriented evidence; B = inconsistent or limited-quality patient-oriented evidence; C = consensus, disease-oriented evidence, usual practice, expert opinion, or case series. For information about the SORT evidence rating system, go to <https://www.aafp.org/afpsort>.

†—Low reactive level laser therapy is the application of light in wavelengths that penetrate the skin, triggering biochemical changes to reduce pain and inflammation. A course of treatment typically involves 30 to 60 seconds of treatment a few times per week for several weeks.

Information from references 5, 10-12, and 64-69.

TABLE 5

**Multi- or Interdisciplinary Pain Rehabilitation for Chronic Musculoskeletal Pain: Review of the Evidence\***

**Common types of pain (SOR rating†)**

Low back	Radicular/referred	Fibromyalgia
<ul style="list-style-type: none"> <li>● Short term: ↓ to ↓↓ pain and ↑ to ↑↑ function (B)<sup>10-12,71</sup></li> <li>● Intermediate term: ↓ pain and ↑ function (B)<sup>10,71</sup></li> <li>● Long term: reports differ (↓ pain and ↑ function vs. no benefit)<sup>10-12,71</sup></li> <li>● Compared with exercise alone, multi- or interdisciplinary pain rehabilitation is better for ↓ pain and ↑ function in the short and intermediate term but not long term (B)<sup>10</sup></li> <li>● Compared with physical therapy alone, multi- or interdisciplinary pain rehabilitation is better for short-term ↓ pain and ↑ function (B)<sup>12</sup></li> <li>● Multi- or interdisciplinary pain rehabilitation ≈ surgery for pain, disability, and work outcomes at two years among candidates for spinal fusion (B)<sup>71</sup></li> </ul>	<ul style="list-style-type: none"> <li>● Insufficient evidence for lumbar radiculopathy<sup>11</sup></li> </ul>	<ul style="list-style-type: none"> <li>● Short term: ↑ function but no pain benefit (B)<sup>10</sup></li> <li>● Intermediate term: ↓ pain and ↑ function (B)<sup>10</sup></li> <li>● Long term: ↑ function but no pain benefit (B)<sup>10</sup></li> <li>● Multi- or interdisciplinary pain rehabilitation ≈ aerobic exercise in improving pain and function in the long term (B)<sup>10</sup></li> </ul>

**Note:** short term = 1 to < 6 months; intermediate term = 6 to < 12 months; long term = ≥ 12 months; ↓ small or undefined decrease; ↑ small or undefined increase; ↓↓ moderate decrease; ↑↑ moderate increase; small = 5 to 10 points on a 100-point scale or SMD of 0.2 to < 0.5; moderate = 10 to 20 points on a 100-point scale or SMD of 0.5 to < 0.8; ● = evidence to support; ● = insufficient or inconsistent evidence; ● = evidence against; ● = comparative effectiveness with other therapies.

SMD = standardized mean difference; SOR = strength of recommendation.

\*—Multi- or interdisciplinary pain rehabilitation includes coordinated care by a team comprised of combinations of clinicians, physical or occupational therapists, and behavior therapists.

†—A = consistent, good-quality patient-oriented evidence; B = inconsistent or limited-quality patient-oriented evidence; C = consensus, disease-oriented evidence, usual practice, expert opinion, or case series. For information about the SORT evidence rating system, go to <https://www.aafp.org/afpsort>.

Information from references 10-12 and 71.

vary from small to no benefit.<sup>10-12,71</sup> There is insufficient evidence regarding the effectiveness of multi- or interdisciplinary pain rehabilitation for lumbar radiculopathy.<sup>11</sup>

A Cochrane review showed that pain, disability, and work outcomes among candidates for spinal fusion were similar at two years between those treated surgically and those treated with multi- or interdisciplinary pain rehabilitation.<sup>71</sup> Multi- or interdisciplinary pain rehabilitation results in short-, intermediate-, and long-term functional improvement and intermediate-term pain reduction in patients with fibromyalgia.<sup>10</sup> The cost-effectiveness of this treatment approach may be enhanced by targeting patients who are most likely to benefit (e.g., those with severe functional deficits, lack of response to less intensive therapies, or significant psychosocial components).<sup>11</sup>

**SUMMARY**

Multi- or interdisciplinary pain rehabilitation appears effective for short- and intermediate-term improvement of low back pain and fibromyalgia, and cost-effectiveness may be enhanced by careful patient selection. The evidence for

multi- or interdisciplinary pain rehabilitation therapy is summarized in *Table 5*.<sup>10-12,71</sup>

**Data Sources:** An Ovid search was completed using the following key terms: low back pain, chronic pain, neck pain, osteoarthritis, fibromyalgia. The following treatment categories were searched: exercise therapy, physical therapy modalities, braces, mind-body therapies, acupuncture therapy, rehabilitation, psychotherapy, musculoskeletal manipulations. Other search terms included noninvasive or non-invasive, nonpharmacologic\* or non-pharmacologic\*; exercise or physical therapy or cognitive or behavioral or feedback or relaxation or acceptance or commitment or traction or ultrasound or stimulation or laser or magnet\* or inferential or electromuscular or diathermy or heat or cold or manipulation or manual or craniosacral or mindfulness or meditation or mind-body or yoga or pilates or Qigong or acupuncture or functional restoration or multidisciplinary\* or interdisciplinary\*. The search included meta-analyses or systematic reviews published in English since January 2018 and studied humans; 2018 was selected as the start date because the AHRQ review included the same search terms through mid-December 2018. Meta-analyses or systematic reviews of trigger point or dry needling\* or imagery (therapies not included in the AHRQ review) were searched for the period 2010 to 2020. Also searched were the Agency for Healthcare Research and Quality Effective

Healthcare Reports, the Cochrane database, and AFP POEMs (Patient-Oriented Evidence that Matters) since 2010. Search date: July 20, 2020.

The views expressed are those of the author and do not reflect the official policy of the Department of the Army, the Department of Defense, or the U.S. government.

### The Author

**DIANE M. FLYNN, MD, MPH**, is the primary care pain management advisor in the Department of Rehabilitative Medicine at the Interdisciplinary Pain Management Center, Madigan Army Medical Center, Tacoma, Wash., and an adjunct assistant professor in the Department of Family Medicine at the Uniformed Services University of the Health Sciences, Bethesda, Md.

Address correspondence to Diane M. Flynn, MD, MPH, Madigan Army Medical Center, 9040 Jackson Ave., Tacoma, WA 98431 (email: [diane.m.flynn4.civ@mail.mil](mailto:diane.m.flynn4.civ@mail.mil)). Reprints are not available from the author.

### References

- Institute of Medicine Committee on Advancing Pain Research, Care, and Education. *Relieving Pain in America: A Blueprint for Transforming Prevention, Care, Education, and Research*. National Academies Press; 2011.
- Murray CJ, Atkinson C, Bhalla K, et al.; U.S. Burden of Disease Collaborators. The state of US health, 1990-2010: burden of diseases, injuries, and risk factors. *JAMA*. 2013;310(6):591-608.
- GBD 2017 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017 [published correction appears in *Lancet*. 2019;393(10190):e44]. *Lancet*. 2018;392(10159):1789-1858.
- Dale R, Stacey B. Multimodal treatment of chronic pain. *Med Clin North Am*. 2016;100(1):55-64.
- Pangarkar SS, Kang DG, Sandbrink F, et al. VA/DoD Clinical practice guideline: diagnosis and treatment of low back pain. *J Gen Intern Med*. 2019;34(11):2620-2629.
- García-Ríos MC, Navarro-Ledesma S, Tapia-Haro RM, et al. Effectiveness of health education in patients with fibromyalgia: a systematic review. *Eur J Phys Rehabil Med*. 2019;55(2):301-313.
- Goodman F, Kaiser L, Kelley C, et al. VA/DoD clinical practice guideline for the non-surgical management of hip and knee osteoarthritis. Department of Veterans Affairs, Department of Defense; 2014. Accessed June 25, 2020. <https://www.healthquality.va.gov/guidelines/cd/oa/index.asp>
- Gross A, Forget M, St George K, et al. Patient education for neck pain. *Cochrane Database Syst Rev*. 2012;(3):CD005106.
- Parreira P, Heymans MW, van Tulder MW, et al. Back schools for chronic non-specific low back pain. *Cochrane Database Syst Rev*. 2017;(8):CD011674.
- Skelly AC, Chou R, Dettori JR, et al. AHRQ Comparative Effectiveness Review, no. 227. Noninvasive nonpharmacological treatment for chronic pain: a systematic review update. April 2020. Accessed July 25, 2020. <https://www.ncbi.nlm.nih.gov/books/NBK556229/>
- Qaseem A, Wilt TJ, McLean RM, et al.; Clinical Guidelines Committee of the American College of Physicians. Noninvasive treatments for acute, subacute, and chronic low back pain: a clinical practice guideline from the American College of Physicians. *Ann Intern Med*. 2017;166(7):514-530.
- Chou R. Nonpharmacologic therapies for low back pain. *Ann Intern Med*. 2017;167(8):606-607.
- Hurley DA, Tully MA, Lonsdale C, et al. Supervised walking in comparison with fitness training for chronic back pain in physiotherapy: results of the SWIFT single-blinded randomized controlled trial (ISRCTN17592092). *Pain*. 2015;156(1):131-147.
- Gross AR, Paquin JP, Dupont G, et al.; Cervical Overview Group. Exercises for mechanical neck disorders: A Cochrane review update. *Man Ther*. 2016;24:25-45.
- Brosseau L, Taki J, Desjardins B, et al. The Ottawa panel clinical practice guidelines for the management of knee osteoarthritis. Part two: strengthening exercise programs. *Clin Rehabil*. 2017;31(5):596-611.
- Fransen M, McConnell S, Harmer AR, et al. Exercise for osteoarthritis of the knee. *Cochrane Database Syst Rev*. 2015;(1):CD004376.
- Brosseau L, Taki J, Desjardins B, et al. The Ottawa panel clinical practice guidelines for the management of knee osteoarthritis. Part three: aerobic exercise programs. *Clin Rehabil*. 2017;31(5):612-624.
- Bartels EM, Juhl CB, Christensen R, et al. Aquatic exercise for the treatment of knee and hip osteoarthritis. *Cochrane Database Syst Rev*. 2016;(3):CD005523.
- Busch AJ, Webber SC, Richards RS, et al. Resistance exercise training for fibromyalgia. *Cochrane Database Syst Rev*. 2013;(12):CD010884.
- Bidonde J, Busch AJ, Schachter CL, et al. Mixed exercise training for adults with fibromyalgia. *Cochrane Database Syst Rev*. 2019;(5):CD013340.
- Bidonde J, Busch AJ, Schachter CL, et al. Aerobic exercise training for adults with fibromyalgia. *Cochrane Database Syst Rev*. 2017;(6):CD012700.
- Bidonde J, Busch AJ, Webber SC, et al. Aquatic exercise training for fibromyalgia. *Cochrane Database Syst Rev*. 2014;(10):CD011336.
- Saragiotto BT, Maher CG, Yamato TP, et al. Motor control exercise for nonspecific low back pain: a Cochrane review. *Spine (Phila Pa 1976)*. 2016;41(16):1284-1295.
- Martin-Gomez C, Sestelo-Diaz R, Carrillo-Sanjuan V, et al. Motor control using cranio-cervical flexion exercises versus other treatments for non-specific chronic neck pain: a systematic review and meta-analysis. *Musculoskeletal Sci Pract*. 2019;42:52-59.
- Tice JA, Kumar V, Otuonye I, et al.; Institute for Clinical and Economic Review. Cognitive and mind-body therapies for chronic low back and neck pain: effectiveness and value. Final evidence report. November 6, 2017. Accessed July 18, 2020. <https://icer-review.org/material/back-and-neck-pain-evidence-report/>
- Nahin RL, Boineau R, Khalsa PS, et al. Evidence-based evaluation of complementary health approaches for pain management in the United States. *Mayo Clin Proc*. 2016;91(9):1292-1306.
- Wieland LS, Skoetz N, Pilkington K, et al. Yoga treatment for chronic non-specific low back pain. *Cochrane Database Syst Rev*. 2017;(1):CD010671.
- Cramer H, Lauche R, Haller H, et al. A systematic review and meta-analysis of yoga for low back pain. *Clin J Pain*. 2013;29(5):450-460.
- Cramer H, Klose P, Brinkhaus B, et al. Effects of yoga on chronic neck pain: a systematic review and meta-analysis. *Clin Rehabil*. 2017;31(11):1457-1465.
- Cramer H, Lauche R, Langhorst J, et al. Yoga for rheumatic diseases: a systematic review. *Rheumatology (Oxford)*. 2013;52(11):2025-2030.
- Kan L, Zhang J, Yang Y, et al. The effects of yoga on pain, mobility, and quality of life in patients with knee osteoarthritis: a systematic review. *Evid Based Complement Alternat Med*. 2016;2016:6016532.
- Langhorst J, Klose P, Dobos GJ, et al. Efficacy and safety of meditative movement therapies in fibromyalgia syndrome: a systematic review and meta-analysis of randomized controlled trials. *Rheumatol Int*. 2013;33(1):193-207.
- Lauche R, Stumpe C, Fehr J, et al. The effects of tai chi and neck exercises in the treatment of chronic nonspecific neck pain: a randomized controlled trial. *J Pain*. 2016;17(9):1013-1027.

## CHRONIC MUSCULOSKELETAL PAIN

34. Ye J, Cai S, Zhong W, et al. Effects of tai chi for patients with knee osteoarthritis: a systematic review. *J Phys Ther Sci*. 2014;26(7):1133-1137.
35. Rendant D, Pach D, Lütke R, et al. Qigong versus exercise versus no therapy for patients with chronic neck pain: a randomized controlled trial. *Spine (Phila Pa 1976)*. 2011;36(6):419-427.
36. Skoglund L, Josephson M, Wahlstedt K, et al. Qigong training and effects on stress, neck-shoulder pain and life quality in a computerised office environment. *Complement Ther Clin Pract*. 2011;17(1):54-57.
37. von Trott P, Wiedemann AM, Lütke R, et al. Qigong and exercise therapy for elderly patients with chronic neck pain (QIBANE): a randomized controlled study. *J Pain*. 2009;10(5):501-508.
38. Geneen LJ, Moore RA, Clarke C, et al. Physical activity and exercise for chronic pain in adults: an overview of Cochrane reviews. *Cochrane Database Syst Rev*. 2017;(4):CD011279.
39. Cherkin DC, Anderson ML, Sherman KJ, et al. Two-year follow-up of a randomized clinical trial of mindfulness-based stress reduction vs. cognitive behavioral therapy or usual care for chronic low back pain. *JAMA*. 2017;317(6):642-644.
40. Monticone M, Cedraschi C, Ambrosini E, et al. Cognitive-behavioural treatment for subacute and chronic neck pain. *Cochrane Database Syst Rev*. 2015;(5):CD010664.
41. Theadom A, Cropley M, Smith HE, et al. Mind and body therapy for fibromyalgia. *Cochrane Database Syst Rev*. 2015;(4):CD001980.
42. Bernardy K, Klose P, Welsch P, et al. Efficacy, acceptability and safety of cognitive behavioural therapies in fibromyalgia syndrome - a systematic review and meta-analysis of randomized controlled trials. *Eur J Pain*. 2018;22(2):242-260.
43. Hempel S, Shekelle PG, Taylor SL, et al. Department of Veterans Affairs. Evidence map of mindfulness. October 2014. Accessed July 18, 2020. [https://www.ncbi.nlm.nih.gov/books/NBK268640/pdf/Bookshelf\\_NBK268640.pdf](https://www.ncbi.nlm.nih.gov/books/NBK268640/pdf/Bookshelf_NBK268640.pdf)
44. Lauche R, Materdey S, Cramer H, et al. Effectiveness of home-based cupping massage compared to progressive muscle relaxation in patients with chronic neck pain—a randomized controlled trial. *PLoS One*. 2013;8(6):e65378.
45. Zech N, Hansen E, Bernardy K, et al. Efficacy, acceptability and safety of guided imagery/hypnosis in fibromyalgia - a systematic review and meta-analysis of randomized controlled trials. *Eur J Pain*. 2017;21(2):217-227.
46. Meeus M, Nijs J, Vanderheiden T, et al. The effect of relaxation therapy on autonomic functioning, symptoms and daily functioning, in patients with chronic fatigue syndrome or fibromyalgia: a systematic review. *Clin Rehabil*. 2015;29(3):221-233.
47. Rubinstein SM, van Middelkoop M, Assendelft WJ, et al. Spinal manipulative therapy for chronic low-back pain. *Cochrane Database Syst Rev*. 2011;(2):CD008112.
48. Coulter ID, Crawford C, Vernon H, et al. Manipulation and mobilization for treating chronic nonspecific neck pain: a systematic review and meta-analysis for an appropriateness panel. *Pain Physician*. 2019;22(2):E55-E70.
49. Gross A, Langevin P, Burnie SJ, et al. Manipulation and mobilisation for neck pain contrasted against an inactive control or another active treatment. *Cochrane Database Syst Rev*. 2015;(9):CD004249.
50. Masaracchio M, Kirker K, States R, et al. Thoracic spine manipulation for the management of mechanical neck pain: a systematic review and meta-analysis. *PLoS One*. 2019;14(2):e0211877.
51. Posadzki P. Is spinal manipulation effective for pain? An overview of systematic reviews. *Pain Med*. 2012;13(6):754-761.
52. Vickers AJ, Vertosick EA, Lewith G, et al.; Acupuncture Trialists' Collaboration. Acupuncture for chronic pain: update of an individual patient data meta-analysis. *J Pain*. 2018;19(5):455-474.
53. Manheimer E, Cheng K, Wieland LS, et al. Acupuncture for hip osteoarthritis. *Cochrane Database Syst Rev*. 2018;(5):CD013010.
54. Deare JC, Zheng Z, Xue CCL, et al. Acupuncture for treating fibromyalgia. *Cochrane Database Syst Rev*. 2013;(5):CD007070.
55. Hu HT, Gao H, Ma RJ, et al. Is dry needling effective for low back pain? A systematic review and PRISMA-compliant meta-analysis. *Medicine (Baltimore)*. 2018;97(26):e11225.
56. Cagnie B, Castelein B, Polлие F, et al. Evidence for the use of ischemic compression and dry needling in the management of trigger points of the upper trapezius in patients with neck pain: a systematic review. *Am J Phys Med Rehabil*. 2015;94(7):573-583.
57. Liu L, Huang QM, Liu QG, et al. Effectiveness of dry needling for myofascial trigger points associated with neck and shoulder pain: a systematic review and meta-analysis. *Arch Phys Med Rehabil*. 2015;96(5):944-955.
58. Furlan AD, Giraldo M, Baskwill A, et al. Massage for low-back pain. *Cochrane Database Syst Rev*. 2015;(9):CD001929.
59. Cherkin DC, Sherman KJ, Kahn J, et al. A comparison of the effects of 2 types of massage and usual care on chronic low back pain: a randomized, controlled trial. *Ann Intern Med*. 2011;155(1):1-9.
60. Perlman AI, Ali A, Njike VY, et al. Massage therapy for osteoarthritis of the knee: a randomized dose-finding trial. *PLoS One*. 2012;7(2):e30248.
61. Salamh P, Cook C, Reiman MP, et al. Treatment effectiveness and fidelity of manual therapy to the knee: a systematic review and meta-analysis. *Musculoskeletal Care*. 2017;15(3):238-248.
62. Li Y, Wang F, Feng C, et al. Massage therapy for fibromyalgia: a systematic review and meta-analysis of randomized controlled trials. *PLoS One*. 2014;9(2):e89304.
63. Liu L, Huang QM, Liu QG, et al. Evidence for dry needling in the management of myofascial trigger points associated with low back pain: a systematic review and meta-analysis. *Arch Phys Med Rehabil*. 2018;99(1):144-152.e2.
64. Gibson W, Wand BM, Meads C, et al. Transcutaneous electrical nerve stimulation (TENS) for chronic pain - an overview of Cochrane reviews. *Cochrane Database Syst Rev*. 2019;(4):CD011890.
65. Martimbianco ALC, Porfirio GJ, Pacheco RL, et al. Transcutaneous electrical nerve stimulation (TENS) for chronic neck pain. *Cochrane Database Syst Rev*. 2019;(12):CD011927.
66. Johnson MI, Claydon LS, Herbison GP, et al. Transcutaneous electrical nerve stimulation (TENS) for fibromyalgia in adults. *Cochrane Database Syst Rev*. 2017;(10):CD012172.
67. Duivenvoorden T, Brouwer RW, van Raaij TM, et al. Braces and orthoses for treating osteoarthritis of the knee. *Cochrane Database Syst Rev*. 2015;(3):CD004020.
68. Yang JD, Tam KW, Huang TW, et al. Intermittent cervical traction for treating neck pain: a meta-analysis of randomized controlled trials. *Spine (Phila Pa 1976)*. 2017;42(13):959-965.
69. Yeh SW, Hong CH, Shih MC, et al. Low-level laser therapy for fibromyalgia: a systematic review and meta-analysis. *Pain Physician*. 2019;22(3):241-254.
70. Jeffery MM, Butler M, Stark A, et al. Agency for Healthcare Research and Quality. Comparative Effectiveness Technical Brief no. 8. Multidisciplinary pain programs for chronic noncancer pain. September 2011. Accessed July 18, 2020. [https://www.ncbi.nlm.nih.gov/books/NBK82511/pdf/Bookshelf\\_NBK82511.pdf](https://www.ncbi.nlm.nih.gov/books/NBK82511/pdf/Bookshelf_NBK82511.pdf)
71. Kamper SJ, Apeldoorn AT, Chiarotto A, et al. Multidisciplinary biopsychosocial rehabilitation for chronic low back pain. *Cochrane Database Syst Rev*. 2014;(9):CD000963.