

Constipation in Children and Adolescents: Evaluation and Treatment

Elie Mulhem, MD, Oakland University William Beaumont School of Medicine, Rochester, Michigan
Faiza Khondoker, DO, and Sanjiv Kandiah, MD, Beaumont Hospital, Troy, Michigan

Constipation in children is usually functional constipation without an organic cause. Organic causes of constipation in children, which include Hirschsprung disease, cystic fibrosis, and spinal cord abnormalities, commonly present with red flag signs and symptoms. A history and physical examination can diagnose functional constipation using the Rome IV diagnostic criteria. The first goal of managing constipation is to treat fecal impaction, and then maintenance therapy is used to prevent a recurrence. Polyethylene glycol is the first-line treatment for constipation. Second-line options include lactulose and enemas. Increasing dietary fiber and fluid intake above usual daily recommendations and adding probiotics provide no additional benefits for treating constipation. Frequent follow-up visits and referrals to a psychologist can assist in reaching some treatment goals. Clinicians should educate caregivers about the chronic course of functional constipation, frequent relapses, and the potential for prolonged therapy. Clinicians should acknowledge caregivers' specific challenges and the negative effects of constipation on the child's quality of life. Referral to a pediatric gastroenterologist is recommended when there is a concern for organic causes or constipation persists despite adequate therapy. (*Am Fam Physician*. 2022;105(5):469-478. Copyright © 2022 American Academy of Family Physicians.)




Illustration by Jennifer Fairman

Constipation in children and adolescents is defined as passing delayed or infrequent hard stools with pain and excessive straining.^{1,2} The prevalence of constipation in children and adolescents is estimated to be as high as 30% worldwide.³ Constipation in children accounts for 3% of primary care physician visits and up to 25% of referrals to pediatric gastroenterologists.¹ Children with constipation incur three times the health care costs of children without constipation,⁴ and chronic constipation can have a negative effect on the child's quality of life.⁵

Normal Defecation Patterns

The number of daily bowel movements decreases as a child ages, averaging four stools per day for infants, two stools per day by two years of age, and one stool per day after four years of age.¹ Infants who are exclusively breastfed are exceptions to these patterns because there may normally be several days between bowel movements.⁶

CME This clinical content conforms to AAFP criteria for CME. See CME Quiz on page 458.

Author disclosure: No relevant financial relationships.

Classification

Functional constipation, for which there is no organic cause, is the most common type of constipation in children and adolescents, accounting for 95% of cases.⁷ Only 5% of constipation cases in children and adolescents can be attributed to an underlying etiology such as Hirschsprung disease, cystic fibrosis, Down syndrome, anorectal malformations, neuromuscular disorders, spinal cord abnormalities, or celiac disease. Certain medications such as opiates, antacids, or anticholinergics can also cause constipation.⁷

Functional constipation is classified as a functional gastrointestinal disorder that cannot be explained by structural or biochemical findings.⁸ Other functional gastrointestinal disorders in children that may be associated with constipation are infant dyschezia and nonretentive fecal incontinence. Infant dyschezia occurs in infants younger than nine months who have constipation-like symptoms, including straining, crying, and transient reddening of the face, but frequently pass soft stools. These symptoms resolve on their own and are usually due to a discoordination of anal sphincter muscles.^{2,7} Nonretentive fecal incontinence differs from functional constipation by the presence of fecal incontinence without stool impaction, a normal number of

SORT: KEY RECOMMENDATIONS FOR PRACTICE

Clinical recommendation	Evidence rating	Comments
Functional constipation is diagnosed by history and physical examination findings in patients without red flag signs or symptoms. ^{6,10}	C	Expert opinion and consensus guideline in the absence of clinical trials
Digital rectal examination and abdominal radiography should not be performed routinely in children with suspected functional constipation. ^{6,12}	C	Expert opinion and consensus guideline in the absence of clinical trials
Increasing fiber or fluid intake above usual daily recommendations does not improve constipation in children. ^{6,24}	A	Consistent evidence from RCTs and systematic reviews
Probiotics are not beneficial in the treatment of constipation in children. ^{6,24,27}	A	Consistent evidence from RCTs and systematic reviews
Referring children with functional constipation to a child psychologist can improve some constipation outcomes. ^{29,30}	B	Inconsistent evidence from RCTs
Polyethylene glycol (Miralax) is first-line treatment for functional constipation in children. ^{5,31,33,34}	A	Consistent evidence from RCTs and systematic reviews

RCT = randomized controlled trials.

A = consistent, good-quality patient-oriented evidence; **B** = inconsistent or limited-quality patient-oriented evidence; **C** = consensus, disease-oriented evidence, usual practice, expert opinion, or case series. For information about the SORT evidence rating system, go to <https://www.aafp.org/afpsort>.

stools, and normal colonic transit time in children four to 18 years of age. Significant psychosocial problems or neurologic lesions may be found in children with nonretentive fecal incontinence.²

Encopresis is not mentioned in the Rome IV diagnostic criteria for constipation; however, it is included in the *Diagnostic and Statistical Manual of Mental Disorders*, 5th ed., as an elimination disorder that can occur with or without fecal retention. Encopresis is the repeated passage of feces in inappropriate places (i.e., clothing or floor), with one or more events occurring each month for three or more months, and the key feature is soiling.^{2,6-10}

Etiology and Pathophysiology

Constipation commonly starts with the transition to solid foods, toilet training, or school entry. The median age of onset of functional constipation is 2.3 years of age.^{2,11} There is no difference in the prevalence of functional constipation between girls and boys.³

Constipation leads to painful bowel movements, which can cause the child to withhold stool. Withholding stool increases colonic water absorption, making the stool firmer and more difficult to pass. The child contracts the anal sphincter or gluteal muscles by stiffening the body to avoid another painful bowel movement. The child may hide in a corner, rock back and forth, or fidget with each urge to defecate. Parents often confuse these withholding behaviors as straining to defecate.¹² Over time, fecal retention stretches

the rectum, which decreases the urge to defecate. The accumulation of stool in the rectum causes a decrease in gastric emptying, resulting in abdominal distention, abdominal pain, nausea, and loss of appetite.² Although children may have painful or traumatic defecation experiences, resulting in stool withholding, adolescents suppress the urge to defecate as a learned behavior.¹³

TABLE 1

Rome IV Diagnostic Criteria for Functional Constipation in Children

≥ 2 of the following, occurring at least once per week for at least 1 month:

- ≤ 2 defecations in the toilet per week in a child ≥ 4 years of age
- ≥ 1 episode of fecal incontinence per week
- History of retentive posturing or excessive volitional stool retention
- History of hard or painful bowel movements
- Presence of large fecal mass in rectum
- Large diameter stools that obstruct toilet

After appropriate medical evaluation, the symptoms cannot be explained by another medical condition

Adapted with permission from Hyams JS, Di Lorenzo C, Saps M, et al. Childhood functional gastrointestinal disorders: children/adolescent. Gastroenterology. 2016;150(6):1464.

Clinical Diagnosis

Functional constipation is a clinical diagnosis based on Rome IV diagnostic criteria (Table 1⁸). The Rome IV criteria are symptom based, and no additional testing is required in patients without red flag signs and symptoms.

A comprehensive history and physical examination (Tables 2 and 3^{2,12,14}) are usually sufficient to diagnose functional constipation and should exclude red flag signs and symptoms suggesting organic causes^{6,10} (Table 4^{9,12,15}). Urinary tract infection symptoms should also be considered in the history because of the possible association with constipation.¹⁶

There is insufficient evidence to support routine digital rectal examination for the diagnosis of constipation, especially in children older than one year with no red flag signs or symptoms.^{6,15} Digital rectal examination may be performed when the diagnosis of constipation is uncertain, to confirm the success of disimpaction treatment, or if organic causes are suspected.

Additional diagnostic testing (i.e., laboratory tests, radiography, colonic transit time, and anorectal manometry) is not required in children with functional constipation and no red flag signs or symptoms.^{6,17,18} Abdominal radiography is of limited value in diagnosing constipation because it does not have interobserver reliability and accuracy. Abdominal radiography should be reserved for specific clinical circumstances in which a rectal examination is needed but undesirable (e.g., in a child with a history of trauma), the diagnosis is uncertain, or to reevaluate following treatment for disimpaction.^{2,6,12}

Differential Diagnosis

The differential diagnosis for children presenting with constipation is listed in Table 5.^{12,17,19} Hirschsprung disease can present with delayed passage of meconium (longer than 48 hours after birth), constipation in the first few months of life, severe abdominal distension, and inadequate weight gain. The physical examination can demonstrate an empty rectum with explosive

stool upon finger withdrawal. Constipation prevalence in children with cystic fibrosis is between 10% to 57%²⁰ and should be considered in patients who have a family history of the disease or present with constipation and other signs such as respiratory symptoms and failure to thrive. Cow's milk protein allergy can cause constipation associated with blood in the stool.²¹ There are also psychological and behavior factors that can result in functional constipation, particularly in patients with autism and attention-deficit/hyperactivity disorders.^{22,23}

TABLE 2

Medical History for the Evaluation of Childhood Constipation

History	Clinical significance
Abdominal pain	It is important to determine whether pain is relieved or affected by defecation (may suggest irritable bowel syndrome); other causes should be ruled out because abdominal pain is often misdiagnosed as being related to constipation
Age at onset of symptoms	Infants younger than 1 month with constipation have a relatively greater likelihood of an organic etiology
Dietary changes	May suggest food allergy or intolerance
Family history, social history, stressors, school entry	May be associated with the onset of constipation
Family's definition of constipation	It is important to assess what caregivers consider as constipation to avoid under/overdiagnosis
Fecal incontinence	Suggests fecal impaction
Hematochezia	May suggest stools that are hard enough to produce fissures or that are associated with an allergy
Other systemic signs (i.e., fever, chills, anorexia, weight loss)	May indicate an organic etiology, such as Hirschsprung disease
Presence of withholding behaviors	Important contributor to constipation in younger children; behavior interventions may be beneficial
Review of current and previous therapies, including diet, behavior, and medications	It is important to understand factors that may influence treatment outcomes
Stool parameters (frequency, consistency using the Bristol stool scale, caliber)	Larger, hard stools may be a sign of withholding; normal bowel movement frequency associated with symptoms may indicate irritable bowel syndrome
Timing of meconium passage after birth	Lack of a bowel movement in first 48 hours suggests Hirschsprung disease
Urinary symptoms	May suggest urinary tract infection secondary to urinary stasis from chronic constipation

Adapted with permission from Nurko S, Zimmerman LA. Evaluation and treatment of constipation in children and adolescents. Am Fam Physician. 2014;90(2):82, with additional information from references 2 and 14.

Treatment

Management of constipation includes dietary modifications, behavior interventions, medications, and disimpaction if needed (Figure 1). Treatment does not differ for preschool-aged children, older children, and adolescents.

DIETARY MODIFICATIONS

Infants respond well to juices containing sorbitol (e.g., apple, prune, pear), which can increase stool frequency and water content in stools. The recommended dose of prune juice for children younger than six months is 1 to 3 mL per kg, which can be diluted with 1 to 2 oz of water.¹² Dietary interventions should be used only as first-line treatment for constipation in infants.¹⁵

TABLE 3

Physical Examination for the Evaluation of Childhood Constipation

Physical examination	Clinical significance
Digital rectal examination (not routinely recommended)	Hirschsprung disease, anorectal malformations
External examination of the perineum and perianal area (look for anal fissures, skin tags)	Hemorrhoids, anorectal malformations
Growth parameters (height, weight, head circumference)	Abnormal growth parameters may indicate an organic etiology (e.g., hypothyroidism, cystic fibrosis)
Lax abdominal musculature	May suggest prune belly (Eagle-Barrett syndrome)
Neurologic evaluation of reflexes (look for cremasteric, anal wink, patellar) and lower extremities (muscle bulk, tone, power)	Spinal cord abnormality
Palpable mass in the lower left quadrant	May suggest impaction
Severe abdominal distension	Obstruction, severe or advanced constipation, Hirschsprung disease
Spine or back examination (look for hair patches, pits, dimples, scoliosis)	Spinal cord abnormality
Thyroid examination (look for thyromegaly, nodules)	Congenital thyroid disease

Information from references 2, 12, and 14.

TABLE 4

Red Flag Signs and Symptoms Suggesting an Organic Cause of Constipation in Children

Signs and symptoms	Potential diagnosis
Abdominal distension	Hirschsprung disease, impaction, neuroenteric condition (e.g., pseudo-obstruction)
Abnormal neurologic examination findings (tight anal sphincter, absent anal wink, absent cremasteric reflex, decreased lower extremity reflexes or tone)	Spinal cord abnormality, anorectal malformation, Hirschsprung disease
Delayed passage of meconium (> 48 hours after birth)	Hirschsprung disease, cystic fibrosis, congenital malformation of anorectum or spine
Extraintestinal symptoms (bilious vomiting, fever, ill appearance)	Hirschsprung disease, neuroenteric condition
Failure to thrive	Hirschsprung disease, cystic fibrosis, metabolic condition, malabsorption
Family history of Hirschsprung disease	Hirschsprung disease
Gushing of stool with rectal examination	Hirschsprung disease
Intermittent diarrhea and explosive stools	Hirschsprung disease
Midline pigmentary abnormalities of lower spine	Spinal cord abnormality
Occult blood in stool	Hirschsprung disease, food allergy, necrotizing enterocolitis
Onset before 1 month of age	Hirschsprung disease, cystic fibrosis, congenital malformation of anorectum or spine, metabolic or endocrine conditions
Perianal fistula, abnormal position of anus	Spinal cord abnormality
Pilonidal dimple covered by tuft of hair	Spinal cord abnormality
Stool caliber	Organic causes such as Hirschsprung disease (constriction from aganglionic segments in Hirschsprung disease results in narrow stools)

Adapted with permission from Nurko S, Zimmerman LA. Evaluation and treatment of constipation in children and adolescents. Am Fam Physician. 2014;90(2):84, with additional information from references 9 and 15.

CONSTIPATION IN CHILDREN

TABLE 5

Differential Diagnosis of Constipation in Children and Recommended Evaluation

Conditions	Recommended diagnostic evaluation (if appropriate)
Anatomic malformations of the colon and rectum	
Anteriorly displaced anus, anal or colonic stenosis, imperforate anus, pelvic mass (sacral teratoma), stricture	Physical examination, barium enema
Drug use/toxin exposure	
Antacids, anticholinergics, antidepressants, antihypertensives, attention-deficit/hyperactivity disorder medications, botulism, calcium channel blockers, chemotherapy, diuretics (dehydration), iron intake, heavy metal toxicity (i.e., lead), opiates, phenobarbital, sucralfate (Carafate), vitamin D toxicity	History, drug level, lead level
Dietary etiologies	
Lack of fiber, starting rice cereal, transitioning from breast milk to formula or cow's milk	Dietary history, cultural patterns of food intake
Functional constipation	History and physical examination; routine testing not needed
Metabolic conditions	
Diabetes insipidus	Serum and urine osmolality
Diabetes mellitus	Fasting glucose level
Hypercalcemia, hypokalemia	Serum calcium and potassium levels
Hypothyroidism	Thyroid studies
Neuropathic gastrointestinal disorders	
Hirschsprung disease	Rectal suction biopsy in children younger than 12 months; full-thickness biopsy for older children
Internal anal sphincter achalasia	Anorectal manometry
Visceral myopathy/neuropathy	Colonic manometry
Neuronal intestinal dysplasia	Anorectal and colonic manometry
Other systemic disorders	
Celiac disease	Tissue transglutaminase IgA, total IgA, small intestine mucosal biopsy
Connective tissue disorder, mitochondrial disorders	Special testing
Cow's milk protein intolerance	Cow's milk elimination from diet
Cystic fibrosis	Sweat test, genetic testing
Psychiatric disorders	Psychological and psychiatric evaluation
Psychosocial	
Birth of sibling, depression/anxiety, parental strife/divorce, starting school, sexual abuse, toilet phobia, toilet training	Social history
Spinal cord anomalies	
Meningocele, spinal cord tumor, tethered cord, trauma	Spinal magnetic resonance imaging, anorectal manometry, urodynamics
Other	
Abnormal abdominal musculature (prune belly [Eagle-Barrett syndrome], gastroschisis, Down syndrome); anismus, Chagas disease, multiple endocrine neoplasia type 2B, porphyria, uremia	As indicated for the suspected condition

IgA = immunoglobulin A.

Adapted with permission from Nurko S, Zimmerman LA. Evaluation and treatment of constipation in children and adolescents. *Am Fam Physician*. 2014;90(2):85, with additional information from references 17 and 19.

CONSTIPATION IN CHILDREN


Although low fluid and fiber intake may contribute to the development of constipation, increasing fluid (water, milk, other beverages) or fiber intake above usual daily recommendations does not improve constipation in children.^{14,24} The recommended daily fluid intake is approximately 4 cups per day for children one to three years of age, 5 cups for children four to eight years of age, and 7 to 8 cups for older children.²⁵ Children and adolescents obtain adequate daily fiber intake from five servings of fruits and vegetables. Recommended daily fiber intake can be calculated by adding 5 g to the child's age in years.²⁶

There is no evidence that increasing physical activity or adding a probiotic supplement alleviates constipation.^{6,27} There is no evidence supporting cow's milk allergy testing for children with constipation.⁶

BEHAVIOR INTERVENTION


Quality of life scores are lower in children with chronic constipation than those without constipation. Children with functional constipation may experience shame, peer

FIGURE 1



Algorithm for the evaluation of constipation in children.

FIGURE 2



Algorithm for the management of functional constipation in children.

CONSTIPATION IN CHILDREN

rejection, bullying, and recurrent treatment failures.^{5,24} Adolescents with functional constipation are at increased risk of depression, anxiety, somatic symptoms, withdrawal, and social problems.²⁸ Current evidence does not support intensive behavior therapy or biofeedback in the treatment of functional constipation. However, minor behavior modifications such as structured toilet training with a reward system, in which the child is instructed to defecate after each meal, may prevent recurrence of fecal impaction.^{6,17}

Some studies have shown that combining medical and behavior therapy can improve fecal incontinence and other clinical outcomes.²⁹⁻³¹ Clinicians should consider referring children with functional constipation to a child psychologist, particularly when a normal defecating pattern is not restored within three months of starting treatment.^{29,30}

Clinicians should acknowledge the possible negative effects of functional constipation on the child's parents or caregivers. Parents caring for children with constipation report having unmet needs, including lack of knowledge about medication use, a sense of isolation, feeling blamed for the child's condition, dealing with treatment failure, and being dismissed by clinicians.³²

DISIMPACTION

For children presenting with fecal impaction, removing the hard stool from the colon is the first step in treating constipation. Fecal impaction can be diagnosed by a history of overflow soiling, the presence of a palpable fecal mass during an abdominal examination, or hard stool in a dilated rectum if a digital rectal examination is warranted.

Disimpaction can be achieved by using oral or rectal medications (*Figure 2*). Polyethylene glycol (Miralax) is a polymer that is minimally absorbed by the gastrointestinal tract, leading to water retention in the intestine and softened stools. Polyethylene glycol is the first-line treatment because it is more effective than other agents for disimpaction and maintenance therapy, is well-tolerated, and has a low risk of adverse effects.^{31,33,34}

For disimpaction, 1 to 1.5 g per kg of polyethylene glycol mixed in 6 to 8 oz of water or juice is given daily for two to six days. Other oral agents are second-line therapy.¹² The North American Society of Pediatric Gastroenterology, Hepatology, and Nutrition recommends using an enema once per day for three to six days as a second-line treatment if polyethylene glycol is not available.⁶

Enemas are not indicated for infants; therefore, glycerin suppositories are often used. Oral and rectal therapies for disimpaction are summarized in *Table 6*.^{2,12}

If disimpaction is not successful at home, hospitalization is required so that polyethylene glycol can be administered by nasogastric tube. This treatment is started at 10 mL per kg per hour and increased by 10 to 20 mL every one to two hours to a maximum dosage of 40 mL per kg per hour. Digital disimpaction is rarely used and is associated with a risk of colon perforation.²

MAINTENANCE THERAPY

Maintenance therapy should be started in children without impaction or when disimpaction is achieved. The

TABLE 6

Therapies for Disimpaction in Children

Therapy	Dosage
Oral	
Polyethylene glycol (Miralax)*	1 to 1.5 g per kg per day
Lactulose	1 mL per kg twice per day for up to 12 weeks
Magnesium citrate	4 mL per kg per day
Rectal	
Enemas (1 per day for 3 to 6 days)	
Saline	5 to 10 mL per kg, administered in the evening
Mineral oil	15 to 30 mL per year of age up to 240 mL
Sodium phosphate	< 2 years: should not be used because of the risk of electrolyte abnormality 2 to 4 years: 1 oz on 2 to 3 consecutive evenings 5 to 11 years: 2.25 oz on 2 to 3 consecutive evenings 12 years: 4.5 oz on 2 to 3 consecutive evenings
Suppository (1 per day)	
Bisacodyl (Dulcolax)	≥ 2 years: 5 to 10 mg (0.5 to 1 suppository)
Glycerin*	0.5 to 1 infant suppository for those up to five years of age; adult suppository for those ≥ 6 years
Other	
Manual (rarely used; general anesthesia needed)	—

*—May be used in infants < 1 year.

Adapted with permission from Nurko S, Zimmerman LA. Evaluation and treatment of constipation in children and adolescents. *Am Fam Physician*. 2014;90(2):87, with additional information from reference 2.

CONSTIPATION IN CHILDREN

goals of maintenance therapy are to achieve soft stool daily or every other day and to prevent stool impaction. Polyethylene glycol is superior to other oral laxatives and is recommended as a first-line maintenance treatment for constipation.^{6,35} The starting maintenance dosage of polyethylene glycol is 0.4 g per kg per day, and caregivers should be advised that it can be increased or decreased to maintain soft stool every day or every other day without diarrhea. Polyethylene glycol appears to be safe for long-term use in children.³⁶ Lactulose is a second-line maintenance treatment if polyethylene glycol is not available. Other second-line therapies include milk of magnesia, stimulant laxatives, and mineral oil.⁶ *Table 7* summarizes

maintenance therapies, including dosing and common adverse effects.^{12,15}

Maintenance therapy should be continued for at least two months or at least one month after a good response to treatment.⁶ Medications should be weaned gradually and may need to be continued for months or years in some patients. If functional constipation develops before toilet training, maintenance therapy should be continued until after toilet training is achieved. Relapses are common and should be treated with maintenance therapy or disimpaction therapy when appropriate.

Rarely, surgical intervention is required to treat refractory constipation. Emerging therapies such as neuromodulation,

TABLE 7

Maintenance Therapies for Children with Constipation

Therapy	Dosage	Adverse effects
First-line osmotics		
Polyethylene glycol (Miralax)*	0.4 to 1.5 g per kg up to 17 g per day	Anaphylaxis, flatulence
Second-line osmotics		
Lactulose*	1 to 3 mL per kg per day in 2 divided doses	Abdominal cramps, flatulence
Milk of magnesia (magnesium hydroxide)	< 2 years: 0.5 mL per kg per day 2 to 5 years: 5 to 15 mL per day 6 to 11 years: 15 to 30 mL per day ≥ 12 years: 30 to 60 mL per day Medication may be given at bedtime or in divided doses	Infants are susceptible to magnesium overdose (hypermagnesemia, hyperphosphatemia, hypocalcemia)
Sorbitol (e.g., prune juice)*	1 to 3 mL per kg once or twice per day in infants	Similar to lactulose
Stimulants		
Bisacodyl (Dulcolax)	3 to 12 years: 5 to 10 mg (1 to 2 tablets) once per day > 12 years: 5 to 15 mg (1 to 3 tablets) once per day	Abdominal cramps, diarrhea, hypokalemia, abnormal rectal mucosa, proctitis (rare), urolithiasis (case reports)
Senna*	1 to 2 years: 1.25 to 2.5 mL (2.2 to 4.4 mg) at bedtime; maximum of 5 mL or 8.8 mg per day 3 to 6 years: 2.5 to 3.75 mL (4.4 to 6.6 mg) or 0.5 tablet (4.3 mg) at bedtime; maximum of 7.5 mL or 1 tablet per day 7 to 12 years: 5 to 7.5 mL (8.8 to 13.2 mg) or 1 tablet (8.6 mg) at bedtime; maximum of 15 mL or 2 tablets per day > 12 years: 10 to 15 mL (26.4 mg) or 2 tablets (17.2 mg) at bedtime; maximum of 30 mL or 4 tablets per day	Idiosyncratic hepatitis, melanosis coli, hypertrophic osteoarthropathy, analgesic nephropathy
Lubricants		
Mineral oil	1 to 3 mL per kg per day divided into 2 doses	Lipoid pneumonia if aspirated, theoretical interference with absorption of fat-soluble substances, foreign body reaction in intestine

*—May be used in infants < 1 year.

Adapted with permission from Nurko S, Zimmerman LA. Evaluation and treatment of constipation in children and adolescents. *Am Fam Physician*. 2014;90(2):88, with additional information from reference 15.

microbiome therapies, and fecal microbiota transplantation may have a role in treating constipation in the future.²⁴ Other organic causes of constipation are treated with the same medications as functional constipation in conjunction with evaluation and treatment of the underlying cause.

Long-term Prognosis

After constipation is diagnosed and the treatment plan is determined, frequent follow-up is recommended to ensure that disimpaction has been achieved and that maintenance therapy is effective.⁷ These follow-up visits also ensure caregiver education and discussions of barriers to implementing treatment recommendations. Clinicians should educate caregivers about functional constipation's chronic course and frequent relapses, and the possible need for prolonged therapy. A patient handout on functional constipation is available at <https://gikids.org/wp-content/uploads/2020/03/Constipation-all-English.pdf>.

A recovery rate of 50% to 60% is reported after one year of intensive treatment,⁶ and about 25% of children continue to have symptoms through adulthood.³⁷ Clinician use of treatment and medication dosing guidelines can decrease the risk of undertreating functional constipation.³⁸ Referral to a pediatric gastroenterologist is recommended if the patient has red flag signs and symptoms or treatment is ineffective.

This article updates previous articles on this topic by Nurko and Zimmerman,¹² and Biggs and Dery.³⁹

Data sources: A PubMed search was completed in Clinical Queries using the key terms constipation, constipated, and children or child. The search included meta-analyses, randomized controlled trials, clinical trials, and reviews. Also searched were Scopus, Embase, Google Scholar, Web of Science, Cochrane database, Essential Evidence Plus, and DynaMed. Search dates: November 19, 2020, and December 14, 2021.

The Authors

ELIE MULHEM, MD, is vice chair and a professor in the Department of Family Medicine and Community Health at Oakland University William Beaumont School of Medicine, Rochester, Mich.

FAIZA KHONDOKER, DO, is a family medicine resident at Beaumont Hospital, Troy, Mich.

SANJIV KANDIAH, MD, is a family medicine resident at Beaumont Hospital.

Address correspondence to Elie Mulhem, MD, Oakland University William Beaumont School of Medicine, 44250 Dequindre Rd., Sterling Heights, MI 48314 (email: emulhem@beaumont.edu). Reprints are not available from the authors.

References

- Madani S, Tsang L, Karnat D. Constipation in children: a practical review. *Pediatr Ann*. 2016;45(5):e189-e196.
- LeLeiko NS, Mayer-Brown S, Cerezo C, et al. Constipation. *Pediatr Rev*. 2020;41(8):379-392.
- Koppen IJN, Vriesman MH, Saps M, et al. Prevalence of functional defecation disorders in children: a systematic review and meta-analysis. *J Pediatr*. 2018;198:121-130.e6.
- Liem O, Harman J, Benninga M, et al. Health utilization and cost impact of childhood constipation in the United States. *J Pediatr*. 2009;154(2):258-262.
- Vriesman MH, Rajindrajith S, Koppen IJN, et al. Quality of life in children with functional constipation: a systematic review and meta-analysis. *J Pediatr*. 2019;214:141-150.
- Tabbers MM, DiLorenzo C, Berger MY, et al.; European Society for Pediatric Gastroenterology, Hepatology, and Nutrition; North American Society for Pediatric Gastroenterology. Evaluation and treatment of functional constipation in infants and children: evidence-based recommendations from ESPGHAN and NASPGHAN. *J Pediatr Gastroenterol Nutr*. 2014;58(2):258-274.
- Khan L. Constipation management in pediatric primary care. *Pediatr Ann*. 2018;47(5):e180-e184.
- Hyams JS, Di Lorenzo C, Saps M, et al. Childhood functional gastrointestinal disorders: children/adolescent. *Gastroenterology*. 2016;150(6):1456-1468.
- Benninga MA, Nurko S, Faure C, et al. Childhood functional gastrointestinal disorders: neonate/toddler. *Gastroenterology*. 2016;150(6):1443-1455.
- Diagnostic and Statistical Manual of Mental Disorders*. 5th ed. American Psychiatric Association; 2014.
- Zeevenhooven J, Koppen IJN, Benninga MA. The new Rome IV criteria for functional gastrointestinal disorders in infants and toddlers. *Pediatr Gastroenterol Hepatol Nutr*. 2017;20(1):1-13.
- Nurko S, Zimmerman LA. Evaluation and treatment of constipation in children and adolescents. *Am Fam Physician*. 2014;90(2):82-90. Accessed November 24, 2021. <https://www.aafp.org/afp/2014/0715/p82.html>
- Youssef NN, Sanders L, Di Lorenzo C. Adolescent constipation: evaluation and management. *Adolesc Med Clin*. 2004;15(1):37-52.
- Waterham M, Kaufman J, Gibb S. Childhood constipation. *Aust Fam Physician*. 2017;46(12):908-912.
- National Institute for Health and Care Excellence. Constipation in children and young people: diagnosis and management. May 26, 2010. Updated July 2017. Accessed December 15, 2021. <https://www.nice.org.uk/guidance/cg99>
- van Summeren JJGT, Holtman GA, van Ommeren SC, et al. Bladder symptoms in children with functional constipation: a systematic review. *J Pediatr Gastroenterol Nutr*. 2018;67(5):552-560.
- Vriesman MH, Koppen IJN, Camilleri M, et al. Management of functional constipation in children and adults. *Nat Rev Gastroenterol Hepatol*. 2020;17(1):21-39.
- Pradhan S, Jagadisan B. Yield and examiner dependence of digital rectal examination in detecting impaction in pediatric functional constipation. *J Pediatr Gastroenterol Nutr*. 2018;67(5):570-575.
- Muise ED, Hardee S, Morotti RA, et al. A comparison of suction and full-thickness rectal biopsy in children. *J Surg Res*. 2016;201(1):149-155.
- Stefano MA, Poderoso RE, Mainz JG, et al. Prevalence of constipation in cystic fibrosis patients: a systematic review of observational studies. *J Pediatr (Rio J)*. 2020;96(6):686-692.
- Miceli Sopo S, Arena R, Greco M, et al. Constipation and cow's milk allergy: a review of the literature. *Int Arch Allergy Immunol*. 2014;164(1):40-45.

CONSTIPATION IN CHILDREN

22. Peeters B, Noens I, Philips EM, et al. Autism spectrum disorders in children with functional defecation disorders. *J Pediatr*. 2013;163(3):873-878.
23. McKeown C, Hisle-Gorman E, Eide M, et al. Association of constipation and fecal incontinence with attention-deficit/hyperactivity disorder. *Pediatrics*. 2013;132(5):e1210-e1215.
24. Southwell BR. Treatment of childhood constipation: a synthesis of systematic reviews and meta-analyses. *Expert Rev Gastroenterol Hepatol*. 2020;14(3):163-174.
25. Rethy J. Choose water for healthy hydration. American Academy of Pediatrics. Updated January 20, 2020. Accessed December 16, 2021. <https://www.healthychildren.org/English/healthy-living/nutrition/Pages/Choose-Water-for-Healthy-Hydration.aspx>
26. Kids need fiber: here's why and how. American Academy of Pediatrics. Updated October 10, 2013. Accessed December 16, 2021. <https://www.healthychildren.org/English/healthy-living/nutrition/Pages/Kids-Need-Fiber-Heres-Why-and-How.aspx>
27. Wojtyniak K, Szajewska H. Systematic review: probiotics for functional constipation in children. *Eur J Pediatr*. 2017;176(9):1155-1162.
28. Rajindrajith S, Ranathunga N, Jayawickrama N, et al. Behavioral and emotional problems in adolescents with constipation and their association with quality of life. *PLoS One*. 2020;15(10):e0239092.
29. Freeman KA, Riley A, Duke DC, et al. Systematic review and meta-analysis of behavioral interventions for fecal incontinence with constipation. *J Pediatr Psychol*. 2014;39(8):887-902.
30. Hankinson JC, Borden L, Allen T, et al. Outcomes of combined medical and behavioral treatments for constipation within a specialty outpatient clinic. *Clin Pract Pediatr Psychol*. 2018;6(1):31-41.
31. Biggs WS, Hamline M, Ebell MH. Constipation (child). Essential Evidence Plus. Updated August 18, 2021. Accessed December 16, 2021. <https://www.essentialevidenceplus.com/content/eee/177>
32. Thompson AP, MacDonald SE, Wine E, et al. Understanding parents' experiences when caring for a child with functional constipation: interpretive description study. *JMIR Pediatr Parent*. 2021;4(1):e24851.
33. Chen SL, Cai SR, Deng L, et al. Efficacy and complications of polyethylene glycols for treatment of constipation in children: a meta-analysis. *Medicine (Baltimore)*. 2014;93(16):e65.
34. Jarzebicka D, Sieczkowska-Golub J, Kierkus J, et al. PEG 3350 versus lactulose for treatment of functional constipation in children: randomized study. *J Pediatr Gastroenterol Nutr*. 2019;68(3):318-324.
35. Gordon M, MacDonald JK, Parker CE, et al. Osmotic and stimulant laxatives for the management of childhood constipation. *Cochrane Database Syst Rev*. 2016;(8):CD009118.
36. Pashankar DS, Uc A, Bishop WP. Polyethylene glycol 3350 without electrolytes: a new safe, effective, and palatable bowel preparation for colonoscopy in children. *J Pediatr*. 2004;144(3):358-362.
37. Bongers ME, van Wijk MP, Reitsma JB, et al. Long-term prognosis for childhood constipation: clinical outcomes in adulthood. *Pediatrics*. 2010;126(1):e156-e162.
38. Borowitz SM, Cox DJ, Kovatchev B, et al. Treatment of childhood constipation by primary care physicians: efficacy and predictors of outcome. *Pediatrics*. 2005;115(4):873-877.
39. Biggs WS, Dery WH. Evaluation and treatment of constipation in infants and children. *Am Fam Physician*. 2006;73(3):469-477. Accessed November 24, 2021. <https://www.aafp.org/afp/2006/0201/p469.html>